

CONFIRM 2F

Supplemental Labeling

SUPPLEMENTAL LABELING FOR USE OF CONFIRM® 2F AGRICULTURAL INSECTICIDE IN FRUITING VEGETABLES

EPA Reg. No 707-238
EPA Est. No. 39578-TX-01

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated. This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of CONFIRM 2F at the recommended rates for a registered use.

Crop	Recropping Interval
Crops for which CONFIRM use is registered.	No restrictions
All other crops	30 days

Note: When using CONFIRM 2F with other registered pesticides, always refer to rotational restrictions and precautions on the other product's label and comply with the most restrictive rotational guidelines.

USE DIRECTIONS FOR FRUITING VEGETABLES*

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult-to-cover crops to insure thorough coverage.

Aerial Application: For optimum performance a minimum application volume of 10 gallons per acre is recommended. Lower carrier volumes may result in less uniform coverage and reduced efficacy.

Spray Adjuvant: One pint of LATRON CS-7 per 100 gallons of spray mixture or similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Alfalfa looper Beet armyworm Black Cutworm Cabbage looper European corn borer Fall armyworm Imported cabbageworm	6.0 to 8.0 (0.09 to 0.12 lb. ai/acre)	For early season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	Do not apply more than 16 fluid ounces per application and do not exceed 64 fluid ounces per season.
Southern armyworm Tobacco hornworm Tomato hornworm True armyworm Yellowstriped armyworm	8.0 to 16.0 (0.12 to 0.25 lb. ai/acre)	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10 to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	Allow at least 7 days to elapse between final application and harvest. See Rotational Crop Restrictions in the body of this labeling.

*The fruiting vegetable grouping includes eggplant, ground cherry, pepino, pepper (bell, chili, cooking), pimiento, tomatillo and tomato.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**



© 2000 Rohm and Haas Company
02/24/00